

The Subscale names on the PACSLAC-II represent the pain assessment domains recommended by the American Geriatrics Society (2002).

Pain Assessment Checklist for Seniors with Limited Ability to Communicate-II (PACSLAC-II)	
Date of Assessment: _____ Time: _____	Check if present
Facial Expressions	
1. Grimacing	
2. Tighter face	
3. Pain expression	
4. Increased eye movement	
5. Wincing	
6. Opening mouth	
7. Creasing forehead	
8. Lowered eyebrows or frowning	
9. Raised cheeks, narrowing of the eyes or squinting	
10. Wrinkled nose and raised upper lip	
11. Eyes closing	
Verbalizations and Vocalizations	
12. Crying	
13. A specific sound for pain (e.g., 'ow', 'ouch')	
14. Moaning and groaning	
15. Grunting	
16. Gasping or breathing loudly	
Body Movements	
17. Flinching or pulling away	
18. Thrashing	
19. Refusing to move	
20. Moving slow	
21. Guarding sore area	
22. Rubbing or holding sore area	
23. Limping	
24. Clenched fist	
25. Going into foetal position	
26. Stiff or rigid	
27. Shaking or trembling	
Changes in Interpersonal Interactions	
28. Not wanting to be touched	
29. Not allowing people near	
Changes in Activity Patterns or Routines	
30. Decreased activity	
Mental Status Changes	
31. Are there mental status changes that are due to pain <u>and</u> are not explained by another condition (e.g., delirium due to medication, etc.)?	
TOTAL SCORE (Add up checkmarks)	

COPYRIGHT: The PACSLAC-II is copyrighted by Sarah Chan, Thomas Hadjistavropoulos and Shannon Fuchs-Lacelle. For permission to reproduce the PACSLAC contact thomas.hadjistavropoulos@uregina.ca who is authorised to provide permission on behalf of all copyright holders. The developers of the PACSLAC-II specifically disclaim any liability arising directly or indirectly from use of application of the PACSLAC-II. Use of the PACSLAC-II may not be appropriate for some patients and the PACSLAC-II is not a substitute for a thorough assessment by a qualified health professional. The PACSLAC-II (like other observational pain assessment tools) is a screening tool and not a definitive indicator of pain. As such, sometimes it may fail to identify pain and other times it may incorrectly signal the presence of pain. The PACSALC-II should be used by qualified health care staff within the context of their broader knowledge and examination of the patient.