

Pain Assessment Checklist for Seniors with Limited Ability to Communicate (PACSLAC)

DATE: _____ TIME ASSESSED: _____

NAME OF PATIENT/RESIDENT: _____

PURPOSE:

This checklist is used to assess pain in patients/residents who have dementia and are unable to communicate verbally.

INSTRUCTIONS:

Indicate with a checkmark, which of the items on the PACSLAC occurred during the period of interest.

Scoring the Sub-Scales is derived by counting the checkmarks in each column. To generate a Total Pain Score sum all four Sub-Scale totals.

Comments:

Facial Expressions	Present
Grimacing	
Sad Look	
Tighter face	
Dirty look	
Change in eyes (squinting, dull, bright, increased movement)	
Frowning	
Pain expression	
Grim face	
Clenching teeth	
Wincing	
Opening mouth	
Creasing forehead	
Screwing up nose	
Activity/Body Movement	
Fidgeting	
Pulling Away	
Flinching	
Restless	
Pacing	
Wandering	
Trying to leave	
Refusing to move	
Thrashing	
Decreased activity	
Refusing medications	
Moving slow	
Impulsive Behaviour (e.g., repetitive movements)	

Activity/Body Movement	Present
Uncooperative/Resistant to care	
Guarding sore area	
Touching/holding sore area	
Limping	
Clenched fist	
Going into foetal position	
Stiff/Rigid	
Social/Personality/Mood	
Physical aggression (e.g., pushing people and/or objects, scratching others, hitting others, striking, kicking)	
Verbal aggression	
Not wanting to be touched	
Not allowing people near	
Angry/Mad	
Throwing things	
Increased confusion	
Anxious	
Upset	
Agitated	
Cranky/Irritable	
Frustrated	
Other*	
Pale Face	
Flushed, red face	
Teary eyed	
Sweating	

Other continued	Present
Shaking/Trembling	
Cold & clammy	
Changes in sleep (please circle):	
Decreased sleep or	
Increased sleep during day	
Changes in appetite (please circle):	
Decreased appetite or	
Increased appetite	
Screaming/Yelling	
Calling out (i.e. for help)	
Crying	
A specific sound or vocalisation For pain 'ow', 'ouch'	
Moaning and groaning	
Mumbling	
Grunting	

Sub-scale Scores:

Facial Expression _____

Activity/Body Movement _____

Social/Personality Mood _____

Other _____

Total Checklist Score _____

* "Other" sub-scale includes physiological changes, eating and sleeping changes and vocal behaviours.

This version of the scale does not include the items "sitting and rocking", "quiet/withdrawn", and "vacant blank stare" as these were not found to be useful in discriminating pain from non-pain states.

Copyright © Shannon Fuchs-Lacelle and Thomas Hadjistavropoulos (The PACSLAC is distributed today with the permission of the copyright holders)

For permission to reproduce the PACSLAC, please contact Thomas Hadjistavropoulos (Thomas.Hadjistavropoulos@uregina.ca)

THE DEVELOPERS OF THE PACSLAC SPECIFICALLY DISCLAIM ANY AND ALL LIABILITY ARISING DIRECTLY OR INDIRECTLY FOR USE OR APPLICATION OF THE PACSLAC. USE OF THE PACSLAC MAY NOT BE APPROPRIATE FOR SOME PATIENTS AND THE PACSLAC IS NOT A SUBSTITUTE FOR A THOROUGH ASSESSMENT OF THE PATIENT BY A QUALIFIED HEALTH PROFESSIONAL. The PACSLAC (like all other related observational tools for seniors with dementia) is a screening tool and not a definitive indicator of pain. As such, sometimes it may incorrectly signal the presence of pain and, other times, it may fail to identify pain. As such, it should be used by qualified health care staff within the context of their broader knowledge and examination of the patient.